



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

or

$$\frac{-1}{4 \sin A \sin B \sin C} \begin{vmatrix} \sin^2 A + \sin^2 B + \sin^2 C, & \sin^2 A + \sin^2 B + \sin^2 C, & \sin^2 A + \sin^2 B + \sin^2 C \\ 1, & 1, & 1 \\ 2 \sin^2 A, & 2 \sin^2 B, & 2 \sin^2 C \end{vmatrix}.$$

But this is equal to zero since two rows are alike after dividing out $\sin^2 A + \sin^2 B + \sin^2 C$. It is to be noticed also that the above determinant is equal to zero for any values whatever of A, B, C provided only that two of them are alike.

Also solved by ELIJAH SWIFT, G. W. HARTWELL, H. POLISH, R. M. MATHEWS, H. L. AGARD, H. S. UHLER, CLIFFORD N. MILLS, W. W. BURTON, CARL A. W. STROM, A. M. KENYON, J. H. WEAVER, and A. H. WILSON.

CALCULUS.

387. Proposed by C. N. SCHMALL, New York City.

Show that the volume bounded by the cone $x^2 + y^2 = (a - z)^2$ and the planes $x = 0, x = z$, is $\frac{2}{3}a^3$.

I. SOLUTION BY A. M. HARDING, University of Arkansas.

The projection on the XY -plane of the curve of intersection of the cone and the plane $z = x$ is $y^2 = a^2 - 2ax$.

If we change this equation to polar coördinates we obtain

$$\rho = \frac{a}{1 + \cos \theta} = \frac{a}{2} \sec^2 \frac{\theta}{2}.$$

Hence,

$$\begin{aligned} \frac{v}{2} &= \int_0^{\pi/2} \int_0^{a/2 \sec^2 \theta/2} \int_x^{a - \sqrt{x^2 + y^2}} dz \cdot \rho d\rho d\theta = \int_0^{\pi/2} \int_0^{a/2 \sec^2 \theta/2} (a - \sqrt{x^2 + y^2} - x) \rho d\rho d\theta \\ &= \int_0^{\pi/2} \int_0^{a/2 \sec^2 \theta/2} (a\rho - \rho^2 - \rho^2 \cos \theta) d\rho d\theta = \frac{a^3}{24} \int_0^{\pi/2} \sec^4 \frac{\theta}{2} d\theta = \frac{a^3}{9}. \end{aligned}$$

Hence the entire volume is $v = \frac{2}{3}a^3$.

II. SOLUTION BY GEO. W. HARTWELL, Hamline University.

If this volume is sliced parallel to the xy plane, the sections between $z = 0$ and $z = a/2$ will be segments of circles and from $z = a/2$ to $z = a$ semicircles.

Integrating then we have

$$\begin{aligned} \int_0^{a/2} \left[\frac{1}{2} \pi (a - z)^2 + z \sqrt{a^2 - 2az} - (a - z)^3 \cos^{-1} \frac{z}{a - z} \right] dz + \int_{a/2}^a \frac{1}{2} \pi (a - z)^2 dz \\ = \int_0^a \frac{1}{2} \pi (a - z)^2 dz + \int_0^{a/2} \left[z \sqrt{a^2 - 2az} - (a - z)^3 \cos^{-1} \frac{z}{a - z} \right] dz. \end{aligned}$$

Integrating we have

$$\begin{aligned} \left[-\frac{1}{6} \pi (a - z)^3 \right]_{z=0}^{z=a} + \left[-\frac{(2a^2 + 6az)(a^2 - 2az)^{3/2}}{30a^2} + \frac{(a - z)^3}{3} \cos^{-1} \frac{z}{a - z} - \frac{a^2 \sqrt{a^2 - 2az}}{3} \right. \\ \left. + \frac{(2a^2 + 2az) \sqrt{a^2 - 2az}}{9} - \frac{(2a^2 + 2az + 3z^2) \sqrt{a^2 - 2az}}{45} \right]_{z=0}^{z=a/2} = \frac{2}{3} a^3. \end{aligned}$$

Also solved by H. L. AGARD, FRANK R. MORRIS, NORMAN ANNING, H. S. UHLER, and the PROPOSER.

388. Proposed by PAUL CAPRON, U. S. Naval Academy.

If $f(x, y) = 0$ represents a curve having a simple tangency to the axis of x at the origin, the